

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

SANONDA HERBICIDE
DIURON 900WG

ACTIVE CONSTITUENT: 900g/Kg DIURON

GROUP C HERBICIDE

*For selective weed control in the situations
as per the Directions for Use table.*

THIS LEAFLET IS PART OF THE LABEL

APVMA Approval No.: 65536/51149



SANONDATM
(AUSTRALIA) PTY LTD
ABN 23 059 813 973

SANONDA (Australia) Pty Ltd
Suite 822, St Kilda Road Towers,
1 Queens Road, Melbourne, VIC 3004
Website: www.sanonda.oz.com

TEL: (03) 9863 8081
FAX: (03) 9863 8083
E-mail: email@sanonda.oz.com

DIRECTIONS FOR USE:**RESTRAINTS:**

DO NOT re-plant treated areas within 2 years of application of Sanonda Herbicide Diuron 900WG except when otherwise stated on label.

DO NOT use on very light sandy or gravelly soils as crop damage may occur.

CONSERVATION TILLAGE

CROP	WEEDS CONTROLLED	STATE	RATE/HA	CRITICAL COMMENTS
Apples & Pears (established at least 1 year)	Barnyard grass, Capeweed, Barley grass, Fat hen, Amaranthus, Mustards, Ryegrass, Wild oats, Wild radish, Wild turnip, Blackberry nightshade, Cobbler's pegs, Crowsfoot grass, Guinea grass, Pigweed, Sowthistle, Summer grass	All States	Single application 4 kg/ha	Apply under trees as a broadcast or band spray. Area should be cleared of weeds prior to spraying Sanonda Herbicide Diuron 900WG. SINGLE APPLICATION: Apply in winter. SPLIT APPLICATION: Apply post harvest, with a second application in spring.
	Wireweed, Toad rush	Qld, NSW, Vic, ACT, SA, WA only	Split application 2 kg/ha	
	Willow herb	Tas only		
Asparagus	Annual Broadleaf weeds and some Annual Grasses	All States	1.5 - 2 kg/ha	Apply in established beds after they have "hilled up" and before emergence of spikes. This will give 10-14 weeks freedom from weeds. Post-emergence application may be made, but the spray should be directed so as to contact only the base of the stems. Use higher rates on heavier soils. DO NOT USE ON SEEDLING ASPARAGUS.
Bananas	Amaranthus, Barley grass, Barnyard grass, Capeweed, Cobbler's pegs, Crowsfoot grass, Fat hen, Guinea grass, Mustard, Pigweed, Prince of Wales feather, Ryegrass, Sowthistle, Summer grass, Wild radish, Wild turnip.	NSW, Qld, WA only	2 - 4 kg/ha	At planting: Apply following fill-in-planting. DO NOT spray banana foliage. Use higher rate on heavier soil types. Apply in 200 to 500 L water/ sprayed ha. Established plantations: Apply avoiding contact with banana plants. Use higher rate on heavier soil types. Add Paraquat 250 if weed growth is above 8 cm high.
			275 g - 500 g/ha Plus 1.2 or 1.6 L/ha Paraquat 250	
Citrus (established at least 1 year)	Seedlings of: Guinea grass, Barnyard grass, Billygoat weed, Wild turnip, Crowsfoot grass, Cobbler's pegs, Wild mustard, Red natal grass, Feather-top, Wild radish, Rhodes grass, Pigweed, Blackberry nightshade, Spiny burr grass, Winter grass, Wild oats, Capeweed, Wireweed, Fat hen, Sowthistle, Summer grass, Barley grass, Ryegrass, Prince of Wales feather, Amaranthus	All States	2 - 4 kg/ha plus 500 mL Surfactant 600 per 100 L spray mixture	Apply only to trees one year or more old. Apply after cultivation in March or April. DO NOT add Surfactant 600 if spray is likely to contact citrus foliage. DO NOT use on very light sands. Apply in up to 4000 L water/ha. Use higher rate on heavier soils and if weeds up to 15 cm in height are present.
	Couch Grass		350 g/100 L	SPOT SPRAYING - Spray to the point of run-off.

CROP	WEEDS CONTROLLED	STATE	RATE/HA	CRITICAL COMMENTS
Coffee	Seedlings of: Summer grass, Barnyard grass, Billygoat weed, Sourgrass	Qld, NSW only	3.9 kg/sprayed ha	Initial application: Destroy all vegetative growth by spraying with Paraquat 250 before applying Sanonda Herbicide Diuron 900WG.
			1 - 1.9 kg/ha plus 600 mL - 1.1 L Paraquat 250 plus Surfactant 600	Subsequent application: Frequency of application will depend on regrowth. Use higher rates if weed growth is dense.
	Couch grass		710 g/200 L	SPOT SPRAYING - Spray to the point of run-off.
Cotton	Annual Broadleaf weeds and some Annual Grasses	Qld, NSW only	1 - 2 kg/ha	<p>Pre-emergence: Apply at planting or within 7 days of planting and prior to cotton emergence. Seed should be planted at least 4 to 5 cm deep and soil compacted over seeds by a normal planter packer wheel. Use high rate on soils high in clay or organic matter content. DO NOT use on light sandy soils as injury to crop may occur. DO NOT make a second pre-emergent application to the same field during the same crop year as injury to the crop may result. Apply in 300 to 450 L water/ha.</p> <p>Post-emergence (lay-by): Use the higher rate on heavier soil types high in clay and/or organic matter. DO NOT use on light sandy soils low in organic matter as injury to the cotton may result. Apply when cotton is at least 30 cm high, or immediately after the last cultivation. If emerged annual weeds (15 cm or less in height) are present add 170 mL 600 g/L Surfactant /100 L water to the spray solution. Apply Sanonda Herbicide Diuron 900WG as a directed spray to cover the area beneath cotton plants and between rows. Avoid spray contact or drift on to cotton plants as injury may result. DO NOT apply more than once per season. DO NOT re-plant treated areas to any crop within one (1) year after last application as injury may result, except cotton, corn or sorghum (grain sorghum only) which may be planted in the spring following treatment. Apply in 300-500 L water/ha.</p>
Gladioli, Tulips, Daffodils, Lilium and Iris	Seedlings of: Summer grass, Barnyard grass, Mustard, Barley grass, Ryegrass, Wild oats, Capeweed, Wild radish, Wild turnip, Blackberry nightshade, Cobbler's pegs	All States	2 kg/ha	Apply in a high volume spray, to moist, bare soil before weeds emerge. Apply after planting but no later than 4 weeks prior to bulb emergence of weeds
	Amaranthus, Annual ryegrass, Barley grass, Barnyard grass, Blackberry nightshade, Capeweed, Cobblers pegs, Crowsfoot grass*, Fat hen, Guinea grass*, Mustards, Pigweed, Redshank, Sow thistles*, Summer grass, Thistles, Wild oats, Wild radish, Wild turnip, Winter grass, Wireweed	Tas only	1.5 - 2.0 kg/ha	<p>DO NOT re-plant treated areas to any crop within one (1) year after treatment..</p> <p>DO NOT use on light sandy or gravelly soils, or soils low in organic matter as it may cause damage to crops, particularly Liliums.</p> <p>Sufficient moisture in the form of rain or irrigation is necessary within 3-4 days of application, to move the product into the weed root-zone. Heavy rains after application may cause severe crop damage.</p> <p>* may require 2.0 kg/ha for control.</p>

CROP	WEEDS CONTROLLED	STATE	RATE/HA	CRITICAL COMMENTS
Grass Seed Crops Established (at least 1 year old) Fescue	Annual Broadleaf weeds and some Annual Grass weeds	SA only	2.8 kg/ha	Apply at weed emergence by boom spray. Use higher rate for heavier soils. Brome grasses will not be controlled.
		Qld, NSW, Vic, ACT & WA only	3 - 3.3 kg/ha	
		All States	1.9 kg/ha	
Phalaris and Cocksfoot				This rate should be used for follow-up sprays in the second year.
Phalaris cultivars Sirolan and Sirosa			1.7 kg/ha	DO NOT use as a tank mix with Paraquat 250
Lucerne (established for 1 year or more)	Seedlings of Capeweed, Wireweed, Chickweed, Summer grass, Docks, Barnyard grass, Thistles, Billygoat weed, Spiny emex, Ryegrass, Barley grass, Shepherds' purse, Fat hen	All States	1 -1.9 kg/ha	Apply when lucerne has been grazed. Weeds should be very small when treated. Use only in areas where lucerne becomes winter dormant. Use higher rate on heavier soil.
	Black nightshade, Wild oats, Wild radish, Winter grass, Amaranthus, Mustard, Wild turnip	Tas only		
Lupins (on light soils only)	Annual ryegrass, Capeweed, Clovers, Medics, Doublegee, Wild radish, Wild turnip and suppression only of Dock, Sorrel, Wireweed, Wild oats and Brome grass	WA only	1.1 kg/ha	Use through boom spray equipment only. Apply in a minimum of 50 L water/ha. Higher volumes of water will give more even application. Must be applied prior to crop emergence. DO NOT use on white or grey sands
Paw Paws (not for use on trees under 9 months old)	Seedlings of Guinea grass, Crowsfoot grass, Barnyard grass, Billygoat weed, Summer grass, Red natal grass, Giant sensitive plant	Qld, NSW, only	4 kg/ha	Avoid contact with plants. Use higher rates on heavier soils.
			2 kg/ha	Apply a second spray if required. DO NOT exceed 6.5 kg/ha per year. Add Paraquat 250 to Sanonda Herbicide Diuron 900WG if weed growth is more than 8 cm high.
Peas	Annual ryegrass, Barley grass, Brome grass, Capeweed, Doublegee, Wild oats, Wild turnip, Wild radish, Mustards	WA only	835 g - 1.2 kg/ha	DO NOT USE ON SANDY SOILS. Use the lower rate pre-sowing in the low rainfall eastern wheat belt and use the higher rate immediately after sowing in the Great Southern Area. Weed control may be increased by mixing with Trifluralin pre-sowing.
Pineapples	Annual Broadleaf weeds and some Annual Grasses	Qld, NSW only	3.9 kg/ha	Note: DO NOT apply more than 8 kg/ha before fruit differentiation of flower bud formation. Apply as a blanket spray after planting and before weed emergence. Ratoon Crop: Apply immediately prior to the first rains on weed free soil. Apply in up to 4000 L water/ha.
			1.9 kg/ha	Follow-up sprays, if necessary, should be made at this lower rate.

CROP	WEEDS CONTROLLED	STATE	RATE/HA	CRITICAL COMMENTS
Sugar Cane (plant and ratoon crops)	Barnyard grass, Crowsfoot grass, Guinea grass, Summer grass, Barley grass, Ryegrass, Wild oats, Cobbler's pegs, Pigweed, Prince of Wales feather, Amaranthus, Fat hen, Mustard, Sowthistle, Capeweed, Wild radish, Wild turnip	Qld, NSW, only	3.9 kg/ha	Pre-emergence of cane and weeds: Soil surface must be moist at time of application. Rain or irrigation is required within ten (10) days of application
			1.9 - 3.9 kg/ha	Post-emergence of cane: As for pre-emergence. Use lower rate if following an application at planting. If weeds at early post-emergence stage addition of Surfactant 600 at 250 mL/100 L of water will enhance activity, but direct spray to below cane leaves.
			275 g/ha - 3.9 kg/ha plus 1.2 L - 1.6 L Paraquat 250	Add to Paraquat 250 to enhance weed control under favourable growing conditions and in open sunny conditions. Sanonda Herbicide Diuron 900WG at rates up to 500 g/ha can be blanket sprayed. Use a directed spray for higher rates of Sanonda Herbicide Diuron 900WG. Complete spray coverage is essential. For grasses and broadleaved weeds up to 5 cm high use a minimum of 250 L spray solution/ha, increase to 350 L/ha for weeds up to 10 cm high. Use a minimum spray volume of 400 L/ha for weeds >10 cm high and for dense mature weeds. Refer to Paraquat label for rates and full directions. Note: Where broadleaved weeds are a serious problem add 2,4-D sodium salt 700g/L DF to the mixture.
Summer Fallows	Potato weed (<i>Heliotropium europaeum</i>)	SA only	275 g/ha	Only effective when in mixture with Paraquat Diquat 250 at 1.6 L/ha (See Paraquat Diquat 250 label)
Tea plantation	Pre-emergence weed control, especially Guinea grass and other grass weeds.	Qld only	1 kg/ha	Use ONLY on tea headlands where soil movement or erosion into cropping areas or waterways is minimal or nil. Apply one boom spray application per year prior to onset of wet season. Avoid contact of spray with the crop. DO NOT use in crop inter-row areas.
Vineyards	Barnyard grass, Crowsfoot grass, Guinea grass, Summer grass, Barley grass, Ryegrass, Wild oats, Cobbler's pegs, Prince of Wales feather, Amaranthus, Fat hen, Mustard, Sowthistle, Capeweed, Wild radish, Wild turnip	All States	2 - 4 kg/ha	Band Treatment to crop rows. Apply to established vines (at least three [3] years old). Apply to moist soil prior to emergence of weeds. DO NOT apply to vines of butt diameter less than 4 cm. Use the lower rate where annual and winter weeds are to be controlled. Avoid spray drift or contact onto fruit or foliage.
	Winter grass, Blackberry nightshade, Thistles	Tas only		
	Annual Broadleaf weeds and some Annual Grasses	All States	85 g/20 m ²	Under racks - apply after harvest.

CROP	WEEDS CONTROLLED	STATE	RATE/HA	CRITICAL COMMENTS	
Wheat, Barley, Triticale, Cereal Rye, Oats	Capeweed, Doublegee, Mustard, Wild Radish [suppression only], Water Weeds [Toads Rush, Crassula]	WA only	275 g - 550 g *Barley to 390 g only	Post planting application only for Full cut sowing systems that leave a relatively smooth surface, eg. standard combine or airseeder. Apply post plant pre-emergent to crop within 3-4 days of planting. Pre plant application only for sowing systems (eg. NoTill or ZeroTill) which leave open slots or furrows, or where seed is sown along the bottom of the furrow. Even the soil surface where harrowed after sowing. Apply within 7 - 0 days pre plant. Soil Types - use the lower rate on sandy soils with low organic carbon or clay content. Caution: DO NOT use on grey or white sands. Crop to be planted at least 35mm deep. Crop tipping and yellowing may occur particularly when heavy rain falls after sowing. Shallow sown crops may be more at risk. Caution: Severe crop damage will occur if herbicides containing paraquat or diquat contact an emerging crop either as a salvage treatment or via spray drift unless rainfall has been exceptionally low.	
	As above plus suppression of Annual Ryegrass, Barley Grass and Silver Grass		550 g/ha plus 500 mL of a 720 g/L metolachlor to improve suppression of Ryegrass		
Wheat, Barley, Triticale, Cereal Rye			275 g - 550 g plus 830 mL/ha of Trifluralin 480	As above. Must be incorporated within 4 hours of application.	
Wheat, Barley, Oats	Capeweed, Charlock, Iceplant, Melilotus, Mustards, Rough Poppy, Saffron thistle, Sheepweed, Three cornered Jack, Wild turnip, Yellow burr weed, Deadnettle	NSW, Vic, ACT & SA only	500 g/ha	Apply when weeds are at the 2-4 leaf stage and soon after the crop is past the 2-5 leaf stage which is normally within 6 weeks of sowing. DO NOT SPRAY LATER THAN EARLY TILLERING STAGE.	
Wheat, Barley, Triticale, Oats	Capeweed, Doublegee, Erodium, London Rocket, Lupin, Mustard, Rapistrum, Wild Radish, Wild turnip, Storksbill, Wild mustard	WA only	200 g/ha plus 400 mL MCPA 500	Apply when the crop has at least 3 to 5 leaves on the main stem. (Zadoks 13-15)	Apply by boom spray or aircraft. Weeds must be in the 2-5 leaf stage. Crop and weeds should be dry at the time of spraying. DO NOT spray if rain is imminent. Undersown sub-clovers may be slightly retarded. DO NOT apply to undersown medics. Apply in 40-70 L water/ha.
			280 g/ha plus 200 mL 2,4-D Amine 625	Apply when crop has at least 4 to 5 leaves on main stem. (Zadoks 14-15)	
Wheat, Barley	Amsinckia (Yellow burr), Capeweed, Deadnettle, Double gee (Three cornered Jack), Mustard, White ironweed (Sheep weed), Wild radish	NSW, Vic & ACT, SA only	280 g/ha plus 500 mL/ha (NSW, Vic, ACT only) 350 mL/ha (SA only) MCPA 500	Apply by boom spray or aircraft when the crop is at the 3-5 leaf stage (NSW, SA only) 5 leaf stage (Vic only). Weeds must be in the 2-5 leaf stage. DO NOT spray if rain is imminent. DO NOT apply to under-sown medic. In NSW, DO NOT use on Durati, hybrid Titan, Kite or Shortim wheat.	
	Soursob	SA only	710 - 980 g/ha	Boom Spray: Sow crop at least 5 cm deep. Apply when crop is in the 2-5 true leaf stage. Use the lower rate on lighter soils.	

140 mm

170 mm

CROP	WEEDS CONTROLLED	STATE	RATE/HA	CRITICAL COMMENTS
Irrigation Channels & Drainage Ditches	Annual and Perennial Grasses and Broadleaf weeds. (Except Nutgrass, Bindweed and Russian knapweed)	SA, WA, Vic, Qld, Tas only	22 kg/ha OR 220 g/100 m	High Volume or Power Spraying. Apply in 2000 L water. Apply during the non-crop season and when channel is not in use. Soil should be moist and preferably bare i.e. after regrading and cleaning channels. Apply prior to expected seasonal rainfall so as to "fix" it into the soil. If 50-100 mm of rain has not fallen after application and it is necessary to use the irrigation channel, fill it with water and allow to stand for 72 hours, then drain off the water in the channel and run to waste.
		NSW only	40 kg/ha maximum	
Rights of Way, Commercial & Industrial Areas	Annual Grasses, Perennial Grasses and Broadleaf weeds. (Except Nutgrass, Johnson grass, Bindweed and Russian knapweed)	All States		If moderate weed growth i.e. greater than 15 cm in height is present at the time of spraying add a knockdown herbicide such as Amitrole T to the mixture. Apply to short growth. Apply just before or during the active growing season. DO NOT treat during the dry season. Time application so that rainfall afterwards will take the chemical down to the root zone for uptake by the roots. For application by Railway Tanker or Boom Spray apply in 400-800 L water/ha. For power spraying or hand spraying equipment, apply the equivalent of at least 2000 L water/ha. Ensure entire area is treated - including bare ground.
			4.5 - 18 kg/ha 18 - 36 kg/ha 27 - 40 kg/ha	Railway Tanker or Boom Sprayer - Initial Treatment: Annual rainfall 0-500 mm Annual rainfall 500-1000 mm Annual rainfall 1000 mm and over. Use higher rates in the ranges shown on soils with high clay or organic matter content and where maximum duration of control is required.
			3.5 - 7 kg/ha 7 - 13.5 kg/ha 9 - 18 kg/ha	Re - Treatment Annual rainfall 0-500 mm. Annual rainfall 500-1000 mm. Annual rainfall 1000 mm and over.
			220 - 890 g/100 L 890 g - 1.8 kg/100 L 1.3 - 2 kg/100 L	High Volume or Power Spraying - Initial Treatment: Annual rainfall 0-500 mm. Annual rainfall 500-1000 mm. Annual rainfall 1000 mm and over.
			180 - 355 g/100 L 355 - 670 g/100 L 450 - 890	Re - Treatment Annual rainfall 0-500 mm. Annual rainfall 500-1000 mm. Annual rainfall 1000 mm and over.
Small Areas			90 g in 15 L knapsack/ 20 m ²	Apply evenly.
Bore Drains	Prickly acacia, Mimosa bush	Qld only	35.5 kg/ha	Block off bore drain 24 hours prior to spraying and allow drain to empty. Spray 1 m wide strip onto mud. Wait 48 - 72 hours before opening drain again. Water MUST by-pass house tanks and desirable tree stands for 7 days after opening drain. DO NOT delve for 2 weeks.
Non-Crop Areas	Sensitive plant (Giant)	WA only	2 - 4 kg/ha or 20 - 40 g/10L	Apply to moist soil.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD:

DO NOT ALLOW ANIMALS TO DRINK WATER FROM TREATED BORE DRAINS FOR 3 DAYS BEFORE SLAUGHTER FOR HUMAN CONSUMPTION.

HARVEST: NOT REQUIRED WHEN USED AS DIRECTED.

GRAZING: NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

Before opening, carefully read Directions for Use, Precautionary Statements, Safety Directions and First Aid Instructions. Application should be made to moist soil before weeds emerge unless otherwise recommended. Sufficient moisture in the form of rain or irrigation is necessary after application to move the chemical into the weed root zone. Add a surfactant where recommended. When spraying emerged weeds, best results are obtained when weeds are actively growing. Thoroughly clean all traces of this product from application equipment immediately after use, otherwise injury to desirable vegetation may result when equipment is used again. Flush tank, pump, hose and boom with several changes of water after removing nozzle tip and screens (clean these parts separately). Keep from contact with fertilizers, insecticides, fungicides and seeds.

MIXING

Partly fill spray tank with water and agitate while adding this product steadily. Continue agitation after filling and during spraying.

To prevent excessive foaming it is recommended to add Foam-Ex at 20 mL/1000 L water prior to adding Sanonda Herbicide Diuron 900WG to the spray tank.

Agitation: An efficient by-pass agitator or means of mechanical agitation should be employed when using this product.

COMPATIBILITY

This product is compatible with Paraquat/Diquat 250 herbicide and 2,4-D Amine, Paraquat 250, MCPA 500. Add the required amount of product when the spray tank containing Sanonda Herbicide Diuron 900WG is nearly full and being continually agitated.

This product is not compatible with Tribunil*, Dimethoate 400, Mataven*, Diclofop-methyl 375 and Bayleton*.

Resistant Weeds Warning**GROUP C HERBICIDE**

Sanonda Herbicide Diuron 900WG is a member of the urea group of herbicides and has the inhibitors of photosynthesis at photosystem II mode of action. For weed resistance management, Sanonda Herbicide Diuron 900WG is a Group C herbicide. Some naturally-occurring weed biotypes resistant to Sanonda Herbicide Diuron 900WG and other Group C herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Sanonda Herbicide Diuron 900WG or other Group C herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Sanonda (Australia) Pty Ltd accepts no liability for any losses that may result from the failure of Sanonda Herbicide Diuron 900WG to control resistant weeds.

WEED CONTROL NON-CROP LAND

Sanonda Herbicide Diuron 900WG is an effective herbicide for the residual control of many annual and perennial grasses and broadleaf weeds on non-crop land such as rail tracks, roadways, firebreaks, industrial sites, fence lines, storage areas, irrigation and drainage ditches where bare ground is desired. Established perennials such as Bindweed, Johnson grass, Nutgrass and Russian Knapweed are not satisfactorily controlled at rates recommended. Duration of control depends on the amount of chemical applied, soil type, rainfall and particular weed species.

SELECTIVE USE IN CROPS

For selective use in crops, it is important to apply recommended rates correctly. Sanonda Herbicide Diuron 900WG is most effective when applied to soil which has been well tilled (clod free) and is free of weeds and trash. Where not recommended for overall application, care should be taken to avoid spray drift onto fruit or foliage. Wetting agents should only be used where specified.

REPLANTING RESTRICTIONS:**Crop Rotations:**

DO NOT replant treated areas to any crop within two (2) years after last spray unless otherwise stated.

Cotton and Lucerne:

DO NOT replant treated areas to any crop within one (1) year after last spray. Except cotton, corn or grain sorghum which may be planted in the spring of the following year.

Gladioli, Tulip, Daffodils, Liliium, Iris:

DO NOT replant treated areas to any crop within one (1) year after treatment.

Pineapples and Sugarcane:

Treated areas can be replanted to pineapple or sugarcane one (1) year after last spray.

140 mm

170 mm

CROP SAFETY

Heavy rains after application of this product may cause severe crop damage, risk must be assumed by the user.

Crop Safety – Apples and Pears:

DO NOT use under dwarf or semi-dwarf trees.

Crop Safety – Grapevines:

DO NOT apply to vines of butt diameter less than 4 cm. On soils low in clay or organic matter, plant injury may occur if heavy rains follow treatment.

DO NOT USE ON LIGHT SANDY OR GRAVELLY SOILS OR SOILS LOW IN ORGANIC MATTER.

AVOID CONTACT OF SPRAYS WITH CROP FOLIAGE OR FRUIT.

Drift Warning:

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF LIVESTOCK

Low hazard to bees. May be applied at any time as recommended in the Directions for Use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT:

DO NOT contaminate streams, rivers or waterways with the chemical or used container.

STORAGE AND DISPOSAL

KEEP OUT OF REACH OF CHILDREN.

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Shake empty bag into container. DO NOT dispose of undiluted chemicals on site. Puncture or shred and bury the empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Avoid contact with eyes and skin. DO NOT inhale dust or spray mist. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre **Phone Australia 131126.**

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from Sanonda (Australia) Pty Ltd at <http://www.sanonda.com>.

140 mm

170 mm

CONDITIONS OF SALE

"Sanonda (Australia) Pty Ltd ('Sanonda') shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Sanonda's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Sanonda has any authority to add to or alter these conditions."


* Other registered trademarks.

**In case of emergency:
Phone 1800 033 498**

Ask for shift supervisor.
Toll free 24 hours.

180 mm

170 mm



SANONDATM
(AUSTRALIA) PTY LTD
ABN 23 059 813 973

SANONDA (Australia) Pty Ltd
Suite 822, St Kilda Road Towers,
1 Queens Road, Melbourne, VIC 3004
Website: www.sanondaos.com

TEL: (03) 9863 8081
FAX: (03) 9863 8083
E-mail: email@sanondaos.com